

Your Post Care Instructions

Hyalase

It is critical that you follow all aftercare instructions below in addition to any advice given by your practitioner to prevent any complications and achieve the best results from your treatment. If the advice given by your practitioner differs from the below, please follow the advice of your practitioner.

If you have any concerns or questions please call us on 0333 242 8528.

Hyalase® administration can result in anaphylaxis (a severe allergic reaction) which is life threatening and requires immediate medical attention. Symptoms of a severe allergic reaction can include shortness of breath, wheezing, coughing, difficulty swallowing, swelling of the tongue, eyelids, lips, hoarseness of the voice, stomach pain, nausea, or diarrhoea.

If you have any of the above symptoms, please report to your nearest Accident and Emergency Department or call 999 for an ambulance.

After the procedure some other common injection-related reactions might occur. These reactions include redness, swelling, pain, itching, bruising and tenderness at the injection site. They have generally been described as mild to moderate and typically resolve spontaneously after a few days after injection. Bruising may occasionally be more significant.

If you experience severe pain or any skin colour changes, such as an area of skin becoming purple or red, or bruising appears abnormal, please contact the clinic immediately.

The below products are recommended by The Cosmetic Skin Clinic to enhance your results, support skin health, and reduce the risk of complications. If the advice given by your practitioner differs from the information below, please follow your practitioner's guidance.

- ZO Sheer Fluid Broad Spectrum SPF 50
- Obagi Rebalance Recovery Cream
- Tebiskin® Cera Boost Face

It's important to have your follow up treatments and to continue with your treatment plan to achieve the best results. Please call us on 0333 242 8528 to book your next treatment.